

Interferon-Gamma (IFN- γ) as a Potential Radio- and Chemo-Protectant

Renee V. Gardner*

LSU Medical Center, New Orleans, Louisiana

Interferons (IFN) have had increasing clinical usage in the treatment of a variety of disorders, at times being used in combination with chemo- or radiotherapy. However, interferons may have inhibitory effects on hematopoietic stem cell proliferation and the effects of this cytokine's use on long-term hematologic function have not been studied. We performed the competitive repopulation assay in the murine system, using cells exposed to irradiation or single-dose chemotherapy with or without concomitant IFN- γ use. IFN- γ alone had no deleterious effects on hematopoietic stem cell productivity. We measured the repopulating ability of exhaustible multilineage precursors that were present at early stages of marrow repopulation after competitive repopulation (30 days). These progenitors were minimally impacted by cyclophosphamide (CTX) with or without IFN- γ . Irradiation (XRT) and CTX alone produced significant repopulating defects in the most primitive hematopoietic stem cell, PHSC. Addition of IFN to either treatment regimen resulted in protection of PHSC, with improved repopulating ability, although the levels of donor marrow reached control levels only when CTX and IFN were used together. The results of multiple use of IFN with chemotherapy must be studied further, but IFN may offer hematologic radio- and chemoprotection, in addition to its antitumor properties in clinical protocols for treatment of cancers. *Am. J. Hematol.* 58:218–223, 1998.

© 1998 Wiley-Liss, Inc.

Key words: interferon; hematopoietic stem cells; radioprotection; chemoprotection

INTRODUCTION

Interferons, potent inflammatory cytokines, are being used in the treatment of an increasing array of clinical disorders. Besides having a direct antiproliferative effect on human cancer cell lines [1], the interferons are known to increase the responsiveness of cellular immune mediators such as natural killer cells or cytotoxic lymphocytes to other cytokines, which stimulate specific antitumor immune responses, whether in an MHC-restricted fashion or not [2,3]. They also promote the activation and aggregation within tumor tissue of these cells [3]. Its modulation of extracellular matrix protein and restoration of normal adhesion molecule interactions has implications for the treatment of disorders such as chronic myelogenous leukemia [4,5].

These varied anti-tumor activities of interferons have led to the use of interferon-gamma (IFN- γ) in a number of clinical settings, and with widely disparate results. It has been used in combination with chemotherapeutic agents in the treatment of colon, small cell lung and metastatic renal cell carcinoma, high-risk cutaneous melanoma, as well as chronic myelogenous leukemia

[6–12]. Because of the demonstrated potentiation of radiation's antiproliferative effects, it has also been given together with radiation in an effort to improve radiation's efficacy in other tumors [13,14]. After transplantation, it has proven useful in the generation of graft-vs.-host (and possibly graft-vs.-leukemia) reactions, which may be beneficial to individuals undergoing autologous marrow transplantation [15–17].

While cytokines have had increasing utility in clinical settings [18–20], the combination of cytokines with chemotherapy may not be an entirely innocuous combination [21] since they may exacerbate an already present marrow reconstitutive defect that may have associated long-term clinical consequences such as residual cytopenias and marrow hypoplasia [22–28]. The long-term hematologic effects of interferon usage with chemotherapy

*Correspondence to: Dr. Renee V. Gardner, LSU Medical Center, Dept. of Pediatrics, 1542 Tulane Avenue, New Orleans LA 70112.

Received for publication 10 February 1997; Accepted 18 March 1998

or radiation, which may also be quite damaging to marrow hematopoietic stem cells, are not known. We employed the competitive repopulation assay [29] to ascertain whether interferon with chemotherapy or sublethal irradiation leads to impairment of the repopulating ability of the most primitive hematopoietic stem cell, which is responsible for life-long marrow stem cell renewal. The results of these experiments are reported.

MATERIALS AND METHODS

Mice

Congenic strains of mice, C57B16/J (B6) and B6-*Hbb^d-Gpi-1^a* (GPI) (Jackson Laboratory, Bar Harbor, ME), were used in all experiments. They were housed and fed under standard conditions. All the mice used were females, aged 8 to 12 weeks old.

B6 and competitor B6-*Hbb^d-Gpi-1^a* mice have different alleles at their *Hbb* and glucose phosphate isomerase loci. B6 mice are homozygous for the *Hbb^s* locus specifying $\alpha_2\beta_2^s$ (single) hemoglobin (Hb), the Hb migrating rapidly on hemoglobin electrophoresis. They also produce Gpi-1^b glucose phosphate isomerase isoenzyme. The B6 congenic strain are homozygous for the *Hbb^d* locus, having $\alpha_2\beta_2^{\text{dmaj}}$ and $\alpha_2\beta_2^{\text{dmin}}$ (diffuse) Hb, migrating at intermediate and slow speeds on electrophoresis. They produce Gpi-1^a isoenzyme, which again has a different migratory or electrophoretic pattern from the B6 isoenzyme. While the mice also differ at the H-1 locus, this difference has been shown to have no significant bearing on transplantation outcome when the two congenic strains are used [30].

Irradiation and Chemotherapy Experiments

Recombinant mouse interferon-gamma, obtained from Genzyme (Cambridge, MA), was used in these experiments. The cytokine had a specific activity of 1×10^7 U/mg and $\geq 95\%$ purity. Mice, 8 to 12 weeks old, were injected with 1 μ g IFN- γ intraperitoneally (i.p.) 24 hr before the administration of a single dose of 200 mg/kg cyclophosphamide intravenously (i.v.) or 500 cGy of irradiation using a ^{137}Cs -irradiator. Cyclophosphamide (CTX) (Sigma, St. Louis, MO), a commonly used anti-tumor agent, has been used extensively by us and has been shown to significantly impair PHSC repopulating ability with just a single dose of 200 mg/kg [20,21], reducing the stem cell pool available for marrow reconstitution. Marrow cell numbers are restored to control or pre-transplant levels by 4 weeks after treatment [20]. Mice were sacrificed after 4 weeks and the marrow from bilateral femurs and tibias was placed in a single-cell suspension in α -medium or DMEM (Gibco, Grand Island, NY), prior to use in competitive repopulation.

Competitive Repopulation

Bone marrow concentration was adjusted to allow 10^6 donor (B6) cells to be mixed with 10^6 competitor (GPI) cells. Equal aliquots of the mixtures were injected into B6 mice that had received 1,100 cGy. Blood was obtained by retroorbital puncture at 30 and 150 days and then separated by density gradient centrifugation to allow separation of lymphocytes for glucose phosphate isomerase electrophoresis. Densitometric analysis permitted determination of the exact percentages of donor marrow.

Measurement of donor cell percentage in recipients at 30 days permits a study of marrow precursors that give rise to both myeloid and lymphoid lineages, but that lack the long-term repopulating ability of PHSC [31]. Precursors measured at this early date are exhaustible, and probably contribute the most to early repopulation of marrow after the radiation-induced myeloblastation for transplant, although some minor additional reconstitutive effort by PHSC at this early date cannot be ruled out entirely. These cells can be separated from PHSC on the basis of limiting dilution and surface antigenic characteristics, and are not identical to CFU-S [32,33]. The PHSC proliferative effort then becomes predominant after 30 days, with PHSC assuming the major reconstitutive role after this early date. Analysis of donor percentages at the later dates measures PHSC repopulating capacity.

Statistical Analyses

Precursor numbers can be referred to as relative repopulating units (RU) and can be estimated by using the following formula [34]:

$$\text{Relative repopulating units (RU)} = (\%) (\text{no. of } 10^5 \text{ untreated competitor marrow cells}) / (100 - \%), \text{ where } \% = \text{percentage donor cell type.}$$

One RU is defined as the repopulating ability of 10^5 untreated competitor marrow cells. If the donor cell population functions normally, 10^5 donor cells should contain approximately 1 RU and repopulate as well as 10^5 untreated competitor marrow cells. If, however, fewer RU are present, it suggests that the number of cells available among donor cells that are capable of marrow reconstitution are fewer than those found in a comparable pool of competitor cells, and repopulating abilities per cell are reduced or both.

Two to three replicates of each experiment were performed. Levels of significance for comparison of results were determined by ANOVA, with results expressed as means \pm standard error of the means.

RESULTS

Measurement of exhaustible multilineage progenitors (EMP) repopulating ability was made 30 days after com-

TABLE I. Results of Competitive Repopulation After Interferon (IFN)- γ and Cyclophosphamide (CTX) or Irradiation at 30 Days*

Treatment groups	% Donor (B6) cells	Total BM cells	CRU/ 10^5 cells	Total CRU/donor	N
Control	52 \pm 3	5.4 $\times 10^7$	1.1	594	20
XRT	20 \pm 3	4.6 $\times 10^7$	0.3	138	16
CTX	44 \pm 2	5.8 $\times 10^7$	0.8	464	25
IFN	52 \pm 1	4.7 $\times 10^7$	1.1	517	6
IFN/CTX	44 \pm 4	5.9 $\times 10^7$	0.8	472	9
IFN/XRT	28 \pm 4	4.9 $\times 10^7$	0.4	196	11

*Data collected 30 days after marrow transplantation are shown. Control groups were untreated. C57B16/J (B6) mice received 500 R (XRT) or 200 mg/kg CTX as a single intravenous (i.v.) dose. IFN- γ was administered 24 hr before either treatment in a dose of 1 μ g given i.p. For competitive repopulation, equal numbers, e.g., 10^6 cells, were used for the bone marrow mixture from competitor (GPI-1A) and donor (B6) animals, and standard aliquots of this mixture were given to lethally irradiated recipients (B6). Results are shown as mean \pm standard error of the means (S.E.M.), while N = the number of mice used for each group. Donor marrow cell counts are shown after treatment; total repopulating units (RU) and concentration of RU are also given. Concentration of RU is represented per 10^5 untreated competitor cells.

petitive repopulation, and results are shown in Table I. The animals in the control group received aliquots of marrow containing equal proportions of donor and competitor cells. As expected, donor cells contributed approximately 50% of the marrow repopulating effort. Interferon alone did not impair EMP repopulating ability, with the single dose used. Radiation, as previously demonstrated, was deleterious to the repopulating ability of EMP; the donor percentage dropped to 20 \pm 3%. In these experiments, EMP were minimally affected by a single dose of CTX at the dose employed of 200 mg/kg, with 44 \pm 4% donor cells seen. Nor did a combination of interferon and CTX significantly damage repopulating ability; the addition of interferon before CTX administration did not lead to any exaggeration of the minimal drop in donor percentage from control levels (to 44 \pm 2%) seen with CTX alone. The addition of interferon prior to irradiation led to an insignificant rise in donor percentage to 28 \pm 4% from 20 \pm 3% after that treatment modality alone.

Calculation of the concentration of repopulating units and total numbers per animal revealed an expected value of about 1 RU/ 10^5 competitor cells for control. The administration of XRT yielded a 2/3 reduction in RU concentration to 0.3, and a decrease in total RU numbers to 23% of control numbers, to 138 and 594. CTX administration resulted in reductions in RU numbers and concentration, but these numbers were approximately 80% of control values. Again, use of IFN- γ before CTX did not greatly alter these results. IFN- γ with XRT led to slight increases in RU, whether in concentration or absolute numbers, with total RU measuring 196; this can be compared to XRT alone, which again decreased RU numbers to 138. No treatment regimen resulted in significant declines in marrow cell numbers.

Effects on Long-Term Hematopoietic Progenitor Cells or PHSC

The long-term repopulating ability of PHSC was measured at 150 days after competitive repopulation. Results are shown in Table II. We have previously demonstrated severe impairment of PHSC repopulating ability after XRT and these results substantiate this effect. Repopulating ability dropped drastically, with the donor percentage being 4% after prior exposure to XRT. These results are statistically significant ($P < 0.001$). Administration of a single dose of CTX resulted in a decline of donor percentage from 50 \pm 4% for the control group to 33 \pm 3% ($P < 0.05$). With preadministration of interferon before CTX, donor percentages increased to control levels, with a value of 47 \pm 4% being recorded, a significant improvement in PHSC repopulating ability ($P < 0.05$) when compared to CTX alone. IFN provided a beneficial effect when given prior to XRT, as well, although not completely protecting PHSC. Donor percentage was increased to 28 \pm 7% from 4 \pm 2% ($P < 0.05$). Interferon itself did not adversely affect PHSC repopulating ability. In fact, donor percentage rose to 65% with just a single dose of IFN- γ .

Exposure of marrow to XRT led to severe drops in RU numbers and concentration to 18 and 0.04/ 10^5 cells (from 540 and 1.0/ 10^5 cells, respectively, for control cells). The use of IFN- γ 24 hr before administration of XRT produced an at least 10-fold increase in these numbers, with RU concentration being 0.4/ 10^5 cells and the total numbers of RU increasing to 196. CTX use alone led to reductions in RU concentration and total numbers. RU numbers were 406/donor, compared again to 540 in control animals. The use of IFN- γ before CTX was restorative, since RU numbers then rose to control levels. IFN- γ alone resulted in increases in absolute numbers

TABLE II. Results of Competitive Repopulation Assay at Days 150 After Interferon (IFN)- γ and Irradiation or Cyclophosphamide (150 Days)[†]

Treatment groups	% Donor (B6) cells	Total BM cells	CRU/10 ⁵ cells	Total CRU/donor	N
Control	50 \pm 4	5.4 \times 10 ¹⁰	1.0	540	21
XRT	4 \pm 2*	4.6 \times 10 ⁷	0.04	18	15
CTX	33 \pm 3*	5.8 \times 10 ⁷	0.7	406	24
IFN	65 \pm 3	4.7 \times 10 ⁷	1.9	893	7
IFN/CTX	47 \pm 4	5.9 \times 10 ⁷	0.9	531	8
IFN/XRT	28 \pm 7*	4.9 \times 10 ⁷	0.4	196	10

[†]Information regarding experimental groups and abbreviations is detailed in Table I. The results are from competitive repopulation experiments analyzed at 150 days. Results are given as means \pm S.E.M., while N represents the number of mice in each group.

*Significant differences ($P < 0.01$ – 0.001) between the indicated group and the controls as determined by ANOVA.

and concentration of RU. Reductions (or increases) in RU number were not proportionate to declines in marrow cell numbers.

DISCUSSION

Interferon's antitumor and inflammatory characteristics have made it a valuable adjunct in treatment protocols for a variety of cancers. However, IFN- γ is an inhibitor of hematopoiesis and has also been implicated recently in the pathogenesis of aplastic anemia [35,36]. Transduction of the IFN- γ gene into human stromal cells resulted in secretion of IFN- γ and resultant inhibition of long-term culture-initiating cell function, in addition to that of more committed progenitors. Whether the long-term hematologic impairment is the result of induced stem or progenitor cell apoptosis or of direct proliferative cell inhibition is not entirely clear, but IFN- γ is responsible for the induction of nitric oxide synthase, which inhibits hematopoiesis in a dose-dependent fashion [37]. The activation of fas antigen by IFN- γ also may potentiate hematopoietic suppression [38].

While the effects of IFN- γ and chemotherapy on the long-term repopulating abilities of hematopoietic stem cells have not been studied previously, hematologic toxicity is a recognized side-effect of many of the regimens utilizing interferons with other anti-tumor agents [7,10,39]. Gallichio and his associates reported increased hematopoietic toxicity with combined IFN- γ and dideoxynucleoside therapy, utilizing in vitro assays [40], although parallel experience has not been noted for IFN- γ with chemotherapy. The ability of interferons to potentiate the toxicity of irradiation in cell lines or tumor tissue is well known [13,14] but again the hematopoietic system has not been studied.

In addition, the full extent of any cytokine's impact on long-term hematopoietic function is not yet known.

Some cytokines such as interleukin-1 (IL-1) and stem cell factor (SCF) have been reported to have radioprotective and/or chemoprotective properties [41–43]. These conclusions have been based on survival data and in vitro assays. In a manner similar to that used with IFN- γ , IL-1 resulted in a 75–90% survival rate in mice after a treatment that otherwise would be 100% fatal [42]. The mechanism of such protection is not entirely clear. SCF has been shown to increase the number of progenitor cells in mice [44–46], while IL-1 may increase aldehyde dehydrogenase content in stem and progenitor cells, promoting resistance to the toxic effects of some cytotoxic agents, such as cyclophosphamide [47].

The concomitant use of cytokines with chemotherapy possibly increases the damage already induced by the cytotoxic drugs, especially since the clinical doses of cytokines are far in excess of their usual physiologic concentrations. This deleterious effect has been demonstrated by Hornung and Longo [25]. In performing murine serial transplantation after exposure of cells to multiple cycles of cyclophosphamide and GM- or G-CSF, they observed significant reductions in the restorative capacity of bone marrow after two to three serial passages when either cytokine was given after chemotherapy administration.

We had fully expected that IFN- γ , given in conjunction with CTX or irradiation, would further exacerbate the previously documented hematopoietic defect seen after these modalities [22–28]. In these experiments, a single dose of CTX and 500 R of XRT were sufficient to cause significant drops in PHSC repopulating ability, although EMP functional impairment was evident only after XRT. Consequently, IFN- γ exerted no or minimal beneficial effects on EMP function, but it did have demonstrable chemoprotective properties for and offered some radioprotection to PHSC, at least with this limited single-dose usage. The addition of IFN- γ after CTX almost completely reversed the hematopoietic defect induced by the chemotherapeutic agent, while repopulating ability was improved 7-fold after XRT.

This ability to protect cells from the deleterious effects of chemotherapy may stem partially from its inhibition of cell cycle progression or of initiation of proliferation of hematopoietic stem cell populations [35,36]. The prevention of cell cycling has been previously associated with stem cell-sparing after chemotherapy, albeit the non-cycling populations have been more committed stem cell populations, such as CFU-S [48,49]. We speculate that IFN prevented exhaustion among the PHSC pool too by maintaining them in a quiescent state when chemotherapy was administered, thus making them less vulnerable to the drug's cytotoxicity. Alternatively, prevention of cycling among more mature progenitors may allow the PHSC pool to reserve its energies, and preserve its num-

bers for self-renewal rather than inducing them to cycle to replenish the numbers of committed precursors.

Selleri et al. [36] have postulated that chronic exposure of hematopoietic stem cells to IFN- γ and other inflammatory cytokines results in the induction of apoptosis and depletion of hematopoietic stem cell pools, a depletion that may be permanent and ultimately result in bone marrow failure. It is conceivable that XRT, more so than single-dose chemotherapy, might induce stromal damage with resultant inflammatory cytokine production, leading to exposure of hematopoietic progenitors to higher and more sustained or continuous levels of IFN- γ . Exacerbation of XRT-induced hematopoietic defect might result from IFN use because such extensive damage could be more difficult to repair.

We have used single-dose IFN- γ to avoid the possible effects of high and repeated exposure of PHSC to the cytokine, and to demonstrate the effects of limited use of IFN- γ with chemotherapy. Multiple doses of IFN with chemotherapy must be tested before any definitive statement about long-term implications of its combined use with chemotherapy can be made. Experiments are being considered to study this question further.

ACKNOWLEDGMENTS

The author thanks Dr. David Harrison for supplying B6 and competitor mice, C.M. Astle for supervision of electrophoretic analyses, Mrs. Bea Stork and Avis Silva for performance of electrophoresis and densitometric readings, and Ms. Jessye Hilliard for secretarial assistance.

REFERENCES

- Hand A, Pelin K, Mattson K, Linnainmaa K: Interferon (IFN)-alpha and IFN-gamma in combination with methotrexate: In vitro sensitivity studies in four human mesothelioma cell lines. *Anticancer Drug* 6:77, 1995.
- Charak BS, Brown EG, Mazumder A: Induction of antitumor effect by treatment with cyclosporine A plus interferon-gamma after chemotherapy: role of cytotoxic cells. *J Immunother Emphasis Tumor Immunol* 17:131, 1995.
- Toliore T, Stravaravdi P, Polyzonis M, Vakalikos J: Natural killer cell activation after interferon administration in patients with metastatic renal cell carcinoma: An ultrastructural and immunohistochemical study. *Eur Urol* 29:252, 1996.
- Bhati R, McGlane PB, Verfaillie CM: Treatment of marrow stroma with interferon- α restores normal β 1 integrin-dependent adhesion of chronic myelogenous leukemia hematopoietic progenitors. Role of MIP-1 α . *J Clin Invest* 96:931, 1995.
- Lagrotta-Candido JM, Villa-Verde DM, Vanderlei FH, Savino W: Extracellular matrix components of the mouse thymus microenvironment. V. Interferon-gamma modulate as thymic epithelial cell/thymocyte interactions via extracellular matrix ligands and receptors. *Cell Immunol* 170:235, 1996.
- Kohara N, Kitaoka F, Komuta K, Yamamoto M, Motojima K, Kanematsu T: Effective treatment of liver metastases from colon cancer with a combination of gamma-interferon and cisplatin chemotherapy: Report of a case. *Surg Today* 25:357, 1995.
- Bitran JD, Green M, Perry M, Hollis DR, Herndon JE: A phase II study of recombinant interferon-gamma following combination chemotherapy for patients with extensive small cell lung cancer. *CALGB. Am J Clin Oncol* 18:67, 1995.
- Lygidakis NJ, Kosmidis P, Ziras N, Parissis J, Kyparidon E: Combined transarterial targeting locoregional immunotherapy-chemotherapy for patients with unresectable hepatocellular carcinoma: A new alternative for an old problem. *J Interferon Cytokine Res* 15:467, 1995.
- Lauta VM: Interferon and chronic myelogenous leukemia. *Med Oncol* 12:41, 1995.
- Mani S, Poo WJ: Single institution experience with recombinant renal cell carcinoma. *Am J Clin Oncol* 19:149, 1996.
- Hamasaki VK, Vokes EE: Interferons and other cytokines in head and neck cancer. *Med Oncol* 12:23, 1995.
- Hussain M, Benedetti J, Smith RE, Rodriguez GI, Schuller D, Ensley J: Evaluation of 96-hour infusion fluorouracil plus cisplatin in combination with alpha interferon for patients with advanced squamous cell carcinoma of the head and neck. *Cancer* 76:1233, 1995.
- Windbichler GH, Hensler E, Widschwendter M, Posch A, Daxenbichler G, Fritsch E, Morth C: Increased radiosensitivity by a combination of 9-cis-retinoic acid and interferon- γ in breast cancer cells. *Gynecol Oncol* 61:387, 1996.
- Vokes EE: The promise of biochemical modulation in combined modality therapy. *Semin Oncol* 2:29, 1994.
- Kennedy MJ, Vogelsang GB, Jones RJ, Farmer ER, Hess AD, Altomonte V, Huelskamp AM, Davidson NE: Phase I trial of interferon gamma to potentiate cyclosporine-induced graft-versus-host disease in women undergoing autologous bone marrow transplantation for breast cancer. *J Clin Oncol* 12:249, 1994.
- Ratanatharathorn V, Uberti J, Karanes C, Lum LG, Abella E, Dan ME, Hussein M, Sensesbrenner LL: Phase I study of alpha-interferon augmentation of cyclosporine-induced graft-versus-host disease in recipients of autologous bone marrow transplantation. *Bone Marrow Transplant* 13:625, 1994.
- Mehta J, Powles R, Singhal S, Tait D, Swansbury J, Treleaven J: Cytokine-mediated immunotherapy with or without donor leukocytes for poor-risk acute myeloid leukemia relapsing after allogeneic bone marrow transplantation. *Bone Marrow Transplant* 16:133, 1995.
- Morstyn G, Sheridan WP: Hematopoietic growth factors in cancer chemotherapy. *Cancer Chemother Biol Response Modif* 14:353, 1993.
- Felner JM: Clinical use of hematopoietic growth factors for control of infections after high-dose chemotherapy. *Ann NY Acad Sci* 730:235, 1994.
- George MJ: The present and future of hematopoietic cytokines in clinical practice. *Stem Cells (Dayton)* 12 (Suppl 1):249, 1994.
- Quesenberry P: Growth factor frenzy. *Exp Hematol* 21:835, 1993.
- Testa NG, Hendry JH, Molineux G: Long-term bone marrow damage in experimental systems and in patients after radiation of chemotherapy. *Anticancer Res* 5:101, 1985.
- Gardner RV, Lerner C, Astle CM, Harrison DE: Assessing permanent damage to primitive hematopoietic stem cells after chemotherapy using the competitive repopulation assay. *Cancer Chemother Pharmacol* 32:450, 1993.
- Neben S, Helman S, Montgomery M, Ferrara J, Mauch P: Hematopoietic stem cell deficit of transplanted bone marrow previously exposed to cytotoxic agents. *Exp Hematol* 21:156, 1993.
- Hornung RL, Longo DL: Hematopoietic stem cell depletion by restorative growth factor regimens during repeated high-dose cyclophosphamide therapy. *Blood* 80:77, 1992.
- Mauch PM, Witsell A, Lamont C, Dawes D, van Os R: Enhancement of bone marrow stem cell damage by administration of G-CSF after chemotherapy. *Exp Hematol* 24:1101, 1996.
- Gardner RV, Astle CM, Harrison DE: Hematopoietic precursor cell

- exhaustion is a cause of proliferative defect in primitive hematopoietic stem cells (PHSC) after chemotherapy. *Exp Hematol* 25:495, 1997.
28. Mauch P, Constone L, Greenberger J, Knospe W, Sullivan J, Liesveld JL, Deeg HJ: Hematopoietic stem cell compartment: Acute and late effects of radiation therapy and chemotherapy. *Int J Radiat Oncol Biol Phys* 31:1319, 1995.
 29. Harrison DE: Competitive repopulation: A new assay for long-term stem cell functional capacity. *Blood* 55:77, 1980.
 30. Harrison DE: A semiquantitative measure of immune responses against erythropoietic stem cell antigens. *Immunogenetics* 261:123, 1987.
 31. Harrison DE, Zhong RK: The same exhaustible multilineage precursor produces both myeloid and lymphoid cells as early as 3–4 weeks after marrow transplantation. *Proc Natl Acad Sci USA* 89:10132, 1992.
 32. Morrison SJ, Weissman IL: The long-term repopulation of a subset of hematopoietic stem cells is deterministic and isolatable by phenotype. *Immunity* 1:661, 1994.
 33. Zhong RK, Aastle CM, Harrison DE: Distinct developmental patterns of short-term and long-term functioning lymphoid and myeloid precursors defined by limiting dilution analysis in vivo. *J Immunol* 157:138, 1996.
 34. Harrison DE, Jordan CT, Zhong RK, Aastle CM: Primitive hematopoietic stem cells: Direct assay of most productive populations by competitive repopulation with simple binomial correlation and covariance calculations. *Exp Hematol* 21:206, 1993.
 35. Selleri C, Sato T, Anderson S, Young NS, Maciejewski JP: Interferon-gamma and tumor necrosis factor-alpha suppress both early and late stages of hematopoiesis and induce programmed cell death. *J Cell Physiol* 165:538, 1995.
 36. Selleri C, Maciejewski JP, Sato T, Young NS: Interferon-gamma constitutively expressed in the stromal microenvironment of human marrow cultures mediates potent hematopoietic inhibition. *Blood* 87:4149, 1996.
 37. Maciejewski JP, Selleri C, Sato T, Cho HJ, Keefer LK, Nathan CF, Young NS: Nitric oxide suppression of human hematopoiesis in vitro. Contribution to inhibitory action of interferon-gamma and tumor necrosis factor-alpha. *J Clin Invest* 96:1085, 1995.
 38. Maciejewski JP, Selleri C, Anderson S, Young NS: Fas antigen expression on CD34⁺ human marrow cells is induced by interferon gamma and tumor necrosis factor alpha and potentiates cytokine-mediated hematopoietic suppression in vitro. *Blood* 85:3183, 1995.
 39. Pujade-Lauraine E, Guastalla JP, Colombo N, Devillier P, Francois E, Fumoleau P, Monnier A, Nooy M, Mignot L, Bugat R, Marques C, Mousseau M, Netter G, Maloisel F, Larbaoui S, Brandely M: Intra-peritoneal recombinant interferon gamma in ovarian cancer patients with residual disease at second-look laparotomy. *J Clin Oncol* 14:343, 1996.
 40. Gallichio VS, Scott KW, Hughes NK, Tse KF, Gaines H, Kirk PR, Birch NJ: Increased hematopoietic toxicity following administration of interferon- α with combination dideoxynucleoside therapy (zidovudine plus ddI) administered in normal mice. *Life Sci* 56:PL 71, 1995.
 41. Neta R: Radioprotection and therapy of radiation injury with cytokines. *Prog Clin Biol Res* 352:471, 1990.
 42. Dalmau SR, Freitas CS, Tabak DG: Interleukin-1 and tumor necrosis factor-alpha as radio- and chemoprotection of bone marrow. *Bone Marrow Transplant* 12:551, 1993.
 43. Zsebo K, Smith K, Hartley C, Greenbaum M, Cooke K, Rich W, McNeice I: Radioprotection of mice by recombinant rat stem cell factor. *Proc Natl Acad Sci USA* 89:9464, 1992.
 44. Bodine DM, Seidel NE, Zsebo KM, Orlic D: In vivo administration of stem cell factor to mice increases the absolute number of pluripotent hematopoietic stem cells. *Blood* 82:445, 1993.
 45. Harrison DE, Zsebo KM, Aastle CM: Splenic primitive hematopoietic stem cell (PHSC) activity is enhanced by steel factor because of PHSC proliferation. *Blood* 83:3146, 1994.
 46. Van Os R, Lamont C, Witsell A: Radioprotection of bone marrow stem cell subsets by interleukin-1 and kit-ligand: Implications for CFU-S as the responsible target cell population. *Exp Hematol* 25:205, 1997.
 47. Moreb J, Zucali JR, Yhang Y, Colvin MO, Gross MA: Role of aldehyde dehydrogenase in the protection of hematopoietic progenitor cells from 4-hydroperoxycyclophosphamide by interleukin 1 α and tumor necrosis factor. *Cancer Res* 52:1770, 1992.
 48. Paukovitz WR, Moser MH, Binder KA, Paukovits JB: The use of haemoregulatory peptides (pEEDCK monomer and dimer) for reduction of cytostatic drug induced haemopoietic damage. *Cancer Treat Rev* 17:347, 1990.
 49. Kriegler AB, Bernardo D, Verschoor SM: Protection of murine bone marrow by dexamethasone during cytotoxic chemotherapy. *Blood* 83:65, 1994.